

Flavonoids lower Alzheimer's A β production via an NF κ B dependent mechanism

Daniel Paris, Venkat Mathura*, Ghania Ait-Ghezala, David Beaulieu-Abdelahad, Nikunj Patel, Corbin Bachmeier, Michael Mullan

Roskamp Institute, 2040 Whitfield Avenue, Sarasota, Florida 34243, USA; Venkat Mathura - Email: vmathura@rfdn.org; *Corresponding author

Received May 28, 2011; Accepted May 29, 2011; Published June 06, 2011

Abstract:

Alzheimer's disease (AD) is characterized by the brain accumulation of A β peptides and by the presence of neurofibrillary tangles. A β is believed to play an important role in AD and it has been shown that certain flavonoids can affect A β production. Recently, it was suggested that the A β lowering properties of flavonoids are mediated by a direct inhibition the β -secretase (BACE-1) activity, the rate limiting enzyme responsible for the production of A β peptides. Western-blots and ELISAs were employed to monitor the impact of flavonoids on amyloid precursor protein processing and A β production. A cell free chemoluminescent assay using human recombinant BACE-1 was used to assess the effect of flavonoids on BACE-1 activity. The effect of flavonoids on NF κ B activation was determined by using a stable NF κ B luciferase reporter cell line. Molecular docking simulations were performed to predict the binding of flavonoids to the BACE-1 catalytic site. Real time quantitative PCR was used to determine the effect of flavonoids on BACE-1 transcription. We show in a cell free assay that flavonoids are only weak inhibitors of BACE-1 activity. Docking simulation studies with different BACE-1 structures also suggest that flavonoids are poor BACE-1 inhibitors as they appear to adopt various docking poses in the active site pocket and have weak docking scores that differ as a function of the BACE-1 structures studied. Moreover, a weak correlation was observed between the effect of flavonoids on A β production *in vitro* and their ability to lower BACE-1 activity suggesting that the A β lowering properties of flavonoids in whole cells are not mediated via direct inhibition of BACE-1 activity. We found however a strong correlation between the inhibition of NF κ B activation by flavonoids and their A β lowering properties suggesting that flavonoids inhibit A β production in whole cells via NF κ B related mechanisms. As NF κ B has been shown to regulate BACE-1 expression, we show that NF κ B lowering flavonoids inhibit BACE-1 transcription in human neuronal SH-SY5Y cells. Altogether, our data suggest that flavonoids inhibit A β and sAPP β production by regulating BACE-1 expression and not by directly inhibiting BACE-1 activity.

Background:

Alzheimer's disease (AD) is a major health concern among the aging population and is the most prevalent form of dementia. While the cause of the disease remains uncertain, the extracellular senile plaques and the intracellular neurofibrillary tangles constitute the two major neuropathological hallmarks present in the brains of AD patients. Neurofibrillary tangles contain hyperphosphorylated microtubule-associated protein tau, while senile plaques contain a core of β -amyloid (A β) peptides. Although the central role of A β remains to be proven in clinical trials, data accumulated during the past two decades place A β peptides and in particular soluble forms of the peptide as being the main molecule triggering the pathological cascade that eventually leads to AD and initiates tau pathology [1]. A β peptides are derived from the cleavage of the β -amyloid precursor protein (APP) by β - and γ -secretases. The major β -secretase is an aspartyl protease termed BACE-1 (β -site APP cleaving enzyme) [2-4]. BACE-1 cleaves APP within the extracellular domain of APP, resulting in the secretion of the large ectodomain (APPs β) and generating a membrane-tethered C-terminal fragment CTF β or C99 which serves as a substrate for γ -secretase [5]. The multimeric γ -secretase complex cleaves at multiple sites within the transmembranous CTF β generating C-terminally heterogeneous A β peptides ranging between 38 to 43 amino-acid residues in length that are secreted [6]. In addition to BACE-1 and γ -secretase, APP can be

cleaved by α -secretase within the A β domain between Lys¹⁶ and Leu¹⁷, releasing APPs α and generating CTF α or C83 which is further cleaved by γ -secretase to generate an N-terminally truncated A β termed p3. Genetic ablation of BACE-1 completely abolishes A β production, establishing BACE-1 as the major neuronal enzyme responsible for initiating the amyloidogenic processing of APP [7].

Current treatments for AD include cholinesterase inhibitors and glutamate antagonists. Although useful, these symptomatic treatments do not stop the disease process or prevent neuronal degeneration. There is an on-going need for the development of new treatments for AD. It has been suggested that a diet rich in polyphenols including flavonoids may have beneficial effects in AD [8]. Flavonoids are plant metabolites that are dietary antioxidant, and it has been hypothesized that this activity may account for their beneficial effects against dementia [9]. The *Ginkgo biloba* extract EGb761 which contains essentially flavonoids (quercetin, kaempferol and isorhamnetin) and terpene lactones (ginkgolides A,B,C and bilobalide) has also been suggested to have positive effects against dementia and AD [10, 11]. Recently, several flavonoids have been shown to regulate A β production and it has been suggested that these compounds act by directly inhibiting BACE-1 activity [12]. As BACE-1 is the rate limiting enzyme responsible for A β production and is considered to be a prime target for AD, we further investigated whether flavonoids can lower A β

production in whole cells by directly inhibiting BACE-1 activity. We tested the effects of different flavonoids on A β production and APP processing using a cell line overexpressing human APP and attempted to correlate the A β lowering activity of the flavonoids with their BACE-1 inhibitory activity. Moreover, we investigated the binding affinity of flavonoids for the BACE-1 catalytic site using thorough docking simulations to determine whether flavonoids hold promise as BACE-1 inhibitors.

Methodology:

Flavonoids:

Daidzein (4',7-Dihydroxyisoflavone, 7-Hydroxy-3-(4-hydroxyphenyl)-4H-1-benzopyran-4-one), 7-Hydroxy-3-(4-hydroxyphenyl)chromone, genistein (4',5,7-Trihydroxyisoflavone, 5,7-Dihydroxy-3-(4-hydroxyphenyl)-4H-1-benzopyran-4-one), luteolin (3',4',5,7-Tetrahydroxyflavone), kaempferol (3,4',5,7-Tetrahydroxyflavone, 3,5,7-Trihydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one), apigenin (4',5,7-Trihydroxyflavone), quercetin (2-(3,4-Dihydroxyphenyl)-3,5,7-trihydroxy-4H-1-benzopyran-4-one, 3,3',4',5,6-Pentahydroxyflavone), α -naphthoflavone (7,8-Benzoflavone), β -naphthoflavone (5,6-Benzoflavone), acacetin (5,7-Dihydroxy-4'-methoxyflavone), taxifolin (3,3',4',5,7-Pentahydroxyflavanone hydrate), PD98059 (2-(2-Amino-3-methoxyphenyl)-4H-1-benzopyran-4-one) were obtained from Sigma Chemicals (MO, USA). Aminogestienin (4'-Amino-6-hydroxyflavone) and baicalein (5,6,7-Trihydroxyflavone) were obtained from Calbiochem (EMD Chemicals, CA, USA).

A β enzyme-linked immunosorbent assay (ELISA):

7W CHO cells stably transfected with human APP751 [13] were maintained in DMEM (ATCC, VA, USA) medium containing 10% fetal bovine serum (Invitrogen, CA, USA), 1X mixture of penicillin/streptomycin/fungizone mixture (Cambrex, ME, USA) and 0.3% geneticin (Invitrogen, CA, USA) as a selecting agent. Cells were cultured in 96-wells culture plates and treated for 24 hours with different doses of flavonoids as indicated in the figure legend. All flavonoids were diluted in DMSO before being exposed to confluent 7W CHO cells so that the final concentration of DMSO in the culture medium was 0.1%. The control wells were treated with 0.1% DMSO. Human A β 1-40 and A β 1-42 were analyzed in the culture medium by using commercially available sandwich ELISAs (Invitrogen, CA) according to the manufacturer's instructions. All experiments were repeated 3-4 times.

Evaluation of APP processing by Western-blot:

The impact of flavonoids on APP processing was evaluated using 7W CHO cells as we previously published [14]. Briefly, confluent 7W CHO cells were treated for 24 hours with 10 and 20 μ M of flavonoids in 24-well plates. Cellular proteins were extracted with 80 μ L of ice-cold M-PER Reagent (Pierce Biotechnology, Rockford, IL, USA) containing 1mM phenylmethanesulfonyl fluoride, 1X of protease cocktail inhibitor (Roche, Inc., USA) and 1mM sodium orthovanadate. Samples were sonicated, denatured by boiling in Laemmli buffer (Bio-Rad, Hercules, CA, USA) and resolved onto 4-20% gradient polyacrylamide gels (Bio-Rad, Hercules, CA, USA). After electrotransferring onto polyvinylidene difluoride membranes, western-blot were immunoprobed with a 1:1000 dilution of an anti-APP C-terminal (751-770) antibody (EMD Biosciences Inc., San Diego, CA, USA), with an anti-actin antibody (Chemicon, Temecula, CA, USA) used as a reference antibody to ensure that equal amount of proteins were electrotransferred. Additionally, sAPP α was detected by Western-blot in the culture medium surrounding 7W CHO cells using the antibody 6E10 (Signet Laboratories Inc., MA, USA) which recognizes amino acids 1-17 of A β and sAPP β was detected in the culture medium using an anti-human sAPP β antibody (Immuno-Biological Laboratories Co. Ltd., Gunma, Japan).

β -secretase activity measurements:

β -secretase activity was measured using human recombinant BACE-1 (Calbiochem, CA, USA) with a commercially available chemoluminescent assay (Discoverix, CA, USA) following the recommendations of the manufacturer. The β -secretase inhibitor IV (BACE IV) inhibitor was used as a positive control in the assay and was purchased from EMD Chemicals (CA, USA). Briefly, BACE-1 enzymatic reactions were carried out for 2 hours at room temperature with 10 ng/ μ L of BACE-1 recombinant enzyme in 96-well plates in a final volume of 100 μ L. Chemoluminescent signals were quantified on a HTS Synergy multiplate reader from Biotek (VT, USA).

NF κ B luciferase activity:

NF κ B activation was quantified using a stable NF κ B luciferase reporter cell

line of HEK293 cells with chromosomal integration of a luciferase reporter construct regulated by 6 copies of the NF κ B response element (Panomics, CA, USA). Cells were grown in DMEM containing 10% serum, 1% penicillin/streptomycin/fungizone and 100 μ g/ml of hygromycin B. Confluent cells were treated with 20 ng/ml of TNF α (Sigma, MO, USA) to induce NF κ B activation and with a dose range of the different flavonoids for 3 hours. Luciferase activity was detected with the Luc-Screen Extended-Glow from Applied Biosystems (CA, USA) and a Synergy HT Biotek chemoluminescent reader (VT, USA) as we previously described [14].

Evaluation of BACE-1 transcription:

Confluent SH-SY5Y cells (ATCC, VA, USA) grown in DMEM/F12 medium supplemented with 10% fetal bovine serum and 1X mixture of penicillin/streptomycin/fungizone mixture were treated with 20 ng/ml of TNF α , or with 20 μ M of apigenin, luteolin, quercetin and daidzein either alone or in combination with 20 ng/ml of TNF α for 30 minutes at 37°C, 5% CO $_2$ (control wells received the same volume of vehicle used to dissolved the flavonoids). After 30 minutes of incubation, RNA was extracted as we previously described [15]. The quality and purity of the RNA obtained were tested on agarose gels and spectrophotometrically at 260 nm and 280 nm. All RNA samples had an A260/280 absorbance ratio between 1.9 and 2.1. Real time quantitative PCR (RT-qPCR) were performed as we previously described [15]. Briefly, to quantify the transcript levels of the BACE-1 gene by RT-qPCR, a protocol using FastStart TaqMan Probes Master Mix (Roche) reaction was performed in duplicates. All reactions contained 2 μ L of cDNA (20 ng), 10 μ L of the 2 \times Master Mix and 0.5 μ L of 20 μ M of each BACE-1 primer (ttcatcaacggctccaact and ctccaggaggtcgcagg), 250 nM of BACE-1 gene specific probe (#04688058001), 500 nM reference (HPRT) gene primer mix (cgtgattagtgatgaaccag and cgagcaagcgttcagctct), 250nM reference HPRT gene probe (# 05046157001) and DEPC-treated water to a final volume of 20 μ L. The reaction protocol started with a 2-min activation step at 50°C, a 10 min template denaturation step at 95°C, followed by 50 cycles of 95°C for 15 sec and 60°C for 20 sec, BACE-1 mRNA fold change (relative to the hypoxanthine-phosphoribosyl-transferase (HPRT) mRNA) was calculated as previously described [15].

Molecular docking study of flavonoids:

Information regarding selected Flavonoids 2D structure was obtained as SMILES notation from Pubchem database (<http://pubchem.ncbi.nlm.nih.gov/>) for: Genistein, Quercetin, Taxifolin, Kaempferol, PD98059, Luteolin, Apigenin, Daidzein, Aminogestienin, alpha- and beta-naphthoflavone. Each of the 2D ligand string was converted into 3D conformers using LigPrep tool and each conformer was further energy minimized by applying OPLS_2005 force field. Low energy conformer for each ligand was retained and subsequently used for docking experiments. We used GLIDE XP (eXtra Precision release 2010, Schrodinger Inc, USA) to perform docking against two different crystal structures of BACE-1 from the PDB files 2B8L and 2QMF. The ligand bound active site pocket in the two crystal structures were identified and a grid map was generated using the centroid of the bound ligand. We used a Vander-Waals radius scaling factor of 1.0 and a partial charge cutoff of 0.25 to soften the potential of non-polar atoms in the BACE pocket and no other explicit constraints were specified. Each docking run samples a variety of poses for the ligand in the BACE active site pocket. An energetic and empirical scoring criterion is used to rank several thousand docked conformations. The scoring algorithm rewards (favorable interactions) or penalizes (unfavorable interactions) the Glide XP score (scoring function) which can be used to further rank the docked pose. A Van der Waals scaling factor of 0.8 for the ligand was used to soften the potential for non-polar ligand atoms. We also performed a post-dock minimization run to obtain favorable conformations and to reduce any strain. We applied a strain correction factor (to account for excessive strain due to conformational restriction upon binding to the pocket) to the Glide XP score whenever the observed strain energy of bound ligand exceeds 4 kcal/mol. A docked pose with a low Glide XP score was considered to be the most favorable state and ranked as the top conformer.

Statistical analysis:

Data were expressed as means \pm S.E.M of n experiments. The statistical significance of the differences between treatment groups was determined by one-way ANOVA and post-hoc comparisons where appropriate using SPSS 12.0.1 for Windows. The half maximal inhibitory concentration (IC $_{50}$) for the different assays was determined using GraphPad Prism 5 (GraphPad Software, Inc., CA, USA).




Figure 1: Dose dependent effects of flavonoids on A β production. The effect of a dose range of flavonoids on A β 1-40 (A) and A β 1-42 (B) production was investigated using 7W CHO cells overexpressing human APP after 24 hours of treatment. ANOVA reveals a significant main effect of the treatments ($P < 0.001$) and doses ($P < 0.001$) on both A β 1-40 and A β 1-42 production. Post-hoc comparisons show significant differences for A β 1-40 production between the control conditions and the β -naphthoflavone ($P < 0.001$), kaempferol ($P < 0.04$), quercetin ($P < 0.001$), aminogenistein ($P < 0.001$), luteolin ($P < 0.001$), apigenin ($P < 0.001$) and acacetin ($P < 0.001$) treatments but no significant differences between the control conditions and baicalein, α -naphthoflavone, daidzein, genistein, PD98059 and taxifolin ($P > 0.05$) treatments. Post-hoc comparisons show significant differences for A β 1-42 production between the control conditions and the β -naphthoflavone ($P < 0.001$), quercetin ($P < 0.001$), aminogenistein ($P < 0.001$), luteolin ($P < 0.001$) and apigenin ($P < 0.008$) but no significant differences between the control conditions and baicalein, α -naphthoflavone, kaempferol, daidzein, genistein, PD98059, taxifolin and acacetin (C). Graph showing a correlation between A β 1-40 and A β 1-42 values obtained following treatments with different doses of the β lowering flavonoids. A strong correlation between A β 1-40 and A β 1-42 values was observed (Pearson correlation coefficient=0.911; $P < 0.001$) showing that flavonoids inhibit A β 1-40 and A β 1-42 with similar potency.




Figure 2: Effects of different flavonoids on APP processing in 7W CHO cells overexpressing human APP. (A) Effect of a 24 hour treatment with different flavonoids (10 and 25 μ M) on sAPP β and sAPP α production by 7W CHO cells. Luteolin, apigenin, β -naphthoflavone, quercetin and kaempferol appear to reduce sAPP β production. (B) Effect of a 24 hour treatment with different flavonoids at a concentration of 50 μ M on sAPP β and sAPP α production by 7W CHO cells. An inhibition of sAPP β production was observed with quercetin, taxifolin, kaempferol, PD98059, luteolin, apigenin and β -naphthoflavone, however genistein, aminogenistein, baicalein and daidzein did not impact sAPP β production. (C) Impact of a 24 hour treatment on APP C-terminal fragments with different flavonoids at a concentration of 20 μ M. A reduction in APP-CTF β production was observed for luteolin, quercetin, apigenin whereas an accumulation of APP-CTF α and APP-CTF β (suggesting an impact on γ -secretase activity) was observed for daidzein, aminogenistein and genistein. Altogether these data suggest that quercetin, taxifolin,

kaempferol, PD98059, luteolin, apigenin and β -naphthoflavone inhibit the β -cleavage of APP.




Figure 3: Effects of flavonoids on BACE-1 activity in a cell free assay. ANOVA reveals a significant main effect of the treatment groups across the doses tested ($P < 0.001$). Post-hoc comparisons show statistical significant BACE-1 activity inhibition across the doses tested for baicalein ($P < 0.001$), apigenin ($P < 0.02$), luteolin ($P < 0.02$), quercetin ($P < 0.001$) and kaempferol ($P < 0.001$) but no significant effect of α -naphthoflavone, β -naphthoflavone, genistein, daidzein, taxifolin, PD98059 and aminogenistein on BACE-1 activity ($P > 0.05$).




Figure 4. (A) Graph showing the docking scores (Observed Glide XP score) correlation for the flavonoids between two crystal structures for BACE-1 (2B8L and 2QMF). The Different flavonoids used in this study are indicated on the graph by the first two letters as bolded here: **Genistein, Quercetin, Taxifolin, Kaempferol, PD98059, Luteolin, Apigenin, Daizein, Aminogeneistein, alpha- and beta-Naphthofalvone**. No significant correlation ($P > 0.1$) was observed between the two scores. In addition no correlation between the different docking scores and the amount of BACE-1 inhibition induced by flavonoids was observed showing that docking simulations do not predict accurately the affinity of flavonoids for the BACE-1 catalytic site. (B) Overlay of several different BACE-1 PDB structures revealing flexible loops (Loop1 and Loop2) in the active site pocket. Different ligand binding sites (using the following PDB structures: 2B8L, 2QMF, 3CIB, 3L5F, 3KMX) in the BACE-1 catalytic site are indicated by space fill dot representation. The active site aspartyl residues (Asp 32 and Asp 228) are represented in red space fill in the left. (C) Docking of kaempferol to BACE-1 structures. Overlay of crystal structure of BACE-1 from PDB file 2B8L (brown ribbon) and 2QMF (green ribbon) along with the top ranked docked pose of kaempferol. Left Side is a zoom of the active site showing differences in the docking orientation of kaempferol in 2B8L (ligand represented by pink sticks) and 2QMF (white sticks). (D) Docking of quercetin to BACE-1 structures. Overlay of crystal structure of BACE-1 from PDB file 2B8L (brown ribbon) and 2QMF (green

ribbon) along with top ranked docked pose of quercetin. Left Side is a zoom of the active site showing differences in the docking orientation of quercetin in 2B8L (ligand represented by pink sticks) and 2QMF (white sticks).

Results:

Effect of flavonoids on $A\beta$, sAPP β and sAPP α production:

The impact of flavonoids on $A\beta$ production was evaluated using 7W CHO cells overexpressing wild-type human APP. Following 24 hours of treatment with a dose range of flavonoids, the amount of human $A\beta$ 1-40 and $A\beta$ 1-42 in the culture media surrounding the 7W CHO cells was measured by ELISAs. A dose dependent inhibition of $A\beta$ 1-40 and $A\beta$ 1-42 production was observed for β -naphthoflavone, aminogenistein, luteolin, apigenin, quercetin, acacetin and PD98059 whereas genistein, α -naphthoflavone, kaempferol baicalein, daidzein and taxifolin were inefficient for the dose range tested (**Figure 1**). The calculated IC_{50} for $A\beta$ 1-40 and $A\beta$ 1-42 were as follows for the most potent flavonoids identified; β -naphthoflavone (0.9 and 1.1 μ M) < aminogenistein (5.7 and 5.8 μ M) < luteolin (6.5 and 8.8 μ M) < apigenin (7.1 and 10.8 μ M) < quercetin (10.2 and 10.2 μ M). Interestingly, a statistically significant correlation was observed between $A\beta$ 1-40 and $A\beta$ 1-42 values showing that active flavonoids are inhibiting $A\beta$ 1-40 and $A\beta$ 1-42 with the same potency (**Figure 1C**).




Figure 5: Graphs representing the correlation between the inhibition of BACE-1 activity observed in a cell free assay and the inhibition of $A\beta$ 1-40 (A) and $A\beta$ 1-42 (B) production in whole cells for the different flavonoids tested. Only a weak correlation was observed between $A\beta$ production and BACE-1 inhibition (Pearson correlation coefficient=0.471; $P < 0.05$).

We next investigated the effects of flavonoids on sAPP β and sAPP α secretion by 7W CHO cells. None of the flavonoids tested appear to stimulate sAPP α production suggesting that these compounds do not stimulate the α -secretase cleavage of APP (**Figure 2**). Among the flavonoids tested, quercetin, kaempferol, luteolin, apigenin and β -naphthoflavone appear to reduce sAPP β production suggesting an inhibition of the β -cleavage of APP (**Figure 2**). We did not observe a significant impact of PD98059 and aminogenistein (at 10 and 25 μ M) on sAPP β production, however a reduction in sAPP β secretion was observed with a 50 μ M dose of PD98059 but not for aminogenistein, daidzein, genistein and aminogenistein (**Figure 2**) suggesting that among the flavonoids tested only luteolin, apigenin, quercetin, β -naphthoflavone, taxifolin, kaempferol and PD98059 impact the β -cleavage of APP. We analyzed the

effect of some of the flavonoids on APP C-terminal fragments and observed that daidzein, aminogenistein and genistein which were unable to inhibit sAPP β production do not inhibit APP-CTF β generation, and to the contrary induce an accumulation of APP-CTF α and CTF β suggesting inhibition of the γ -secretase cut (Figure 2). A decreased APP-CTF β production as well as an inhibition of sAPP β secretion was observed for luteolin, quercetin, apigenin and kaempferol (Figure 2) suggesting that these flavonoids lower A β production by impacting the β -cleavage of APP.




Figure 6: (A) Dose dependent inhibition of NFkB activity by A β lowering flavonoids. HEK293 cells stably expressing an NFkB luciferase reporter construct were co-treated with 20 ng/ml of TNF α and different doses of the flavonoids for 3 hours before measuring NFkB luciferase activity. ANOVA reveals a significant main effect of the flavonoid treatments ($P < 0.001$) and of the doses used ($P < 0.001$) on NFkB activity. Post-hoc comparisons show statistically significant differences between the control conditions and the treatments with kaempferol, quercetin, luteolin, apigenin and acacetin ($P < 0.001$) on NFkB activity. Graphs representing the correlation between the inhibition of NFkB activity and A β 1-40 (B) and A β 1-42 (C) production observed for the different flavonoids tested. A strong correlation between A β values and the amount of NFkB inhibition induced by flavonoids was observed (Pearson correlation coefficient=0.872; $P < 0.001$).




Figure 7: Effect of daidzein, apigenin, luteolin and quercetin on BACE-1 transcription in human neuronal SH-SY5Y cells. BACE-1 transcription was quantified by RT-PCR in SHSY cells following treatment with flavonoids for 30 minutes in presence and absence of 20 ng/ml of TNF α . A strong induction of BACE-1 mRNA was observed in SH-SY5Y cells challenged with the NFkB inducer TNF α and an approximate 10 fold reduction in BACE-1 transcription was observed following treatments with apigenin, luteolin and quercetin showing that these flavonoids can mitigate BACE-1 expression induced by NFkB stimulation. No inhibition of TNF α induced BACE-1 transcription was observed with daidzein. ANOVA reveals a statistically significant main effect of TNF α ($P < 0.001$), apigenin ($P < 0.001$), luteolin ($P < 0.001$) and quercetin ($P < 0.001$) on BACE-1 mRNA level but no significant main effect of daidzein ($P = 0.323$). Post-hoc comparisons show significant differences between TNF α and TNF α +apigenin ($P < 0.001$), TNF α +luteolin ($P < 0.001$) or TNF α +quercetin treatments ($P < 0.001$) but no statistically significant difference between TNF α and TNF α +daidzein ($P = 0.805$) treatments on BACE-1 mRNA level.

Effects of flavonoids on BACE-1 activity:

As many of the flavonoids tested appear to inhibit the β -cleavage of APP, we determined whether these flavonoids were direct inhibitors of the β -secretase (BACE-1) using a chemoluminescent cell free assay. We selected a chemoluminescent assay over a fluorescent based assay for measuring BACE-1 activity since flavonoids are fluorescent (data not shown) and may interfere with the detection of BACE-1 activity by fluorescence. Among the flavonoids tested, baicalein appears the most potent for inhibiting BACE-1 activity with an IC₅₀ around 10 μ M (Figure 3), however this compound is unable to significantly lower A β or sAPP β production in whole cells (Figure 2). Luteolin, quercetin, kaempferol and apigenin dose dependently inhibited BACE-1 activity (Figure 3). Overall, the flavonoids tested appear weak inhibitors of BACE-1 activity in the cell free assay and the calculated IC₅₀ values were as follows; baicalein (10.2 μ M)<kaempferol (27.7 μ M)<quercetin (30.2 μ M)<apigenin (34.2 μ M)<luteolin (56.5 μ M). To further evaluate the possible interaction of the flavonoids with BACE-1, we performed *in silico* docking simulations. The results of our docking experiments along with the Glide XP score is summarized in Table 1. We observed that the independent docking experiments based on two X-ray structures gave different rank orders of the flavonoids tested. As such, we observed a correlation coefficient of 0.33 ($p = 0.12$) showing no significant relationship between the Glide XP score of the docked flavonoids from the two different docking experiments (Figure 4). We further aligned the docked structure of flavonoids from the different docking experiments and found that the docked poses do not align. We observed that the highest docking score difference was for apigenin which had a Glide XP score of -9.01 while docking to 2B8L and -3.18 while docking to 2QMF. The best score among flavonoids observed while docking to 2B8L was for taxifolin (XP score -10.49) and worst for daidzein (XP score -2.91). While docking flavonoids to 2QMF, we found that the best score was observed for baicalein (-8.81) and the worst for apigenin (-3.18). Based on the observed BACE-1 IC₅₀ in a cell free assay, we found that the docking experiment with 2QMF had a better score for active flavonoids that inhibited BACE-1 at sub-micromolar range. Overall the flavonoids tested appear to be only weak BACE-1 inhibitors. In addition, only a weak correlation between the inhibition of BACE-1 activity and the inhibition of A β production by flavonoids in whole cells (Figure 5) was observed and their IC₅₀ for inhibiting BACE-1 activity in a cell free assay is higher than their IC₅₀ for inhibiting A β production in whole

cells suggesting these compounds mitigate A β production via additional mechanisms independently of direct BACE-1 inhibition.

Effect of flavonoids on NF κ B activity and BACE-1 transcription:

We next investigated the effect of flavonoids on NF κ B activity since flavonoids are known to display anti-inflammatory properties [16] and since we have shown previously that compounds of unrelated structure that inhibit NF κ B activation can lower A β production by targeting the β -cleavage of APP [17, 18]. We observed that all of the flavonoids unable to significantly lower A β level in whole cells for the dose range tested (PD98059, taxifolin, genistein, daidzein and baicalein) were also unable to reduce NF κ B activation induced by TNF α , whereas A β and APPs β lowering flavonoids (kaempferol, quercetin, acacetin, apigenin and luteolin) dose dependently inhibited NF κ B activation (Figure 6). Interestingly, we found a strong correlation between the amount of NF κ B inhibition and the level of A β 1-40 and A β 1-42 inhibition for the different doses of the flavonoids tested, suggesting the A β lowering activity of the flavonoids is mediated via NF κ B (Figure 6). We next investigated the possible impact of daidzein, apigenin, luteolin and quercetin on BACE-1 transcription using human neuronal SHSY cells and confirmed that NF κ B lowering flavonoids (apigenin, luteolin and quercetin) inhibit BACE-1 transcription stimulated by TNF α whereas daidzein, which does not significantly inhibit NF κ B activity, does not affect BACE-1 transcription (Figure 7).

Discussion:

Several studies have revealed that natural flavonoids can reduce A β neurotoxicity [19-21] possibly via an antioxidative mechanism and inhibition of A β oligomerization [22]. In addition, several flavonoids have been shown to lower brain A β accumulation in transgenic mouse models of AD [23, 24]. Flavonoids have been proposed to act as BACE-1 inhibitors, the rate limiting enzyme responsible for the production of A β peptides [12, 25]. In our study, some of the A β lowering flavonoids that we tested decreased sAPP β production suggesting an inhibition of the β -cleavage of APP. However, these flavonoids only weakly inhibited BACE-1 activity in a cell free assay and we found only a weak correlation between their effect on BACE-1 activity and their ability to lower A β in whole cells suggesting their A β lowering properties are mediated principally via other mechanisms. We also investigated the binding of flavonoids to the BACE-1 catalytic site employing docking simulations. Careful mining of the Protein Data Bank (PDB) for BACE-1 revealed that there are 144 entries including apo and complexed structures with several different inhibitors. When we analyzed these structures using a molecular visualization tool, we observed that the active site of BACE-1 possesses a high conformational flexibility upon ligand binding. The active site consists of aspartyl residues (32 and 228) encompassed by two highly flexible loop regions (Loop1: 9-13 KSGQG and Loop2: 71-74 YTQG) that undergo both side chain and backbone rearrangements. For example, in the structure 2B8L the SER10 and its corresponding residue SER71 in the structure 2QMFA, the distance between the backbone atom C- α is 6.83 Å whereas the distance between the side chain atom O- γ is 7.68 Å. The distance between 2B8L:72THR and 3KMXA:133THR for CG2 atom is 5.74 Å and between the backbone C- α atoms is 4.38 Å. The active site pocket is plastic and different structures can be aligned to understand conformational changes due to induced fitting of the ligand. BACE-1 possesses a large active site pocket typical of aspartyl proteases and further adopts conformational changes upon ligand binding that may include both side-chain rearrangement and backbone movement of residues in the active site [26-28]. Due to this fact, docking studies do not adequately reproduce ligand binding and may possibly over or under rank a ligand upon docking resulting in false positives. In our study, we used two different structures for BACE-1 and performed multiple docking studies. We observed that as a function of the different structures selected for BACE-1, the docking score/rank for the flavanoids was different. As a matter of fact, we found no correlation between the observed IC₅₀ for BACE-1 inhibition in the cell-free assay and the Glide XP Score in the docking experiments. Energetic calculations from our BACE-1 docking studies also reveal that the flavonoids only weakly bind the BACE-1 catalytic site; in agreement with the weak inhibition of BACE-1 activity we observed with flavonoids in a BACE-1 cell free assay. Docking studies of BACE-1 do not adequately reproduce the native structure when ligands are cross-docked and hence further studies that combine docking with Monte Carlo/Molecular Dynamics methods may be required [29]. Our data suggest the A β lowering flavonoids we tested are only poor BACE-1 inhibitor and must therefore regulate A β production in whole cells by other mechanisms. This conclusion is different from past studies [12] in which docking results on a single BACE-1

structure were presented as evidence to support the hypothesis that flavonoids directly inhibit BACE-1 activity explaining their A β lowering properties in whole cells. Using docking experiments with two different crystal structures for BACE-1, we demonstrate that the docking score of flavonoids is different for the two BACE-1 structures tested and do not correlate with the inhibition of BACE-1 activity observed with flavonoids in a cell free assay. It is therefore inaccurate to classify active from inactive flavonoids as BACE-1 inhibitors by simply using docking simulations alone. The A β lowering activity of flavonoids cannot be explained solely by their effect on BACE-1 activity. We found a lack of correlation between the amount of BACE-1 inhibition observed and the ability of the compounds to lower A β production in whole cells. In addition, their IC₅₀ for inhibiting BACE-1 activity in a cell free assay was higher than their IC₅₀ for inhibiting A β in whole cells suggesting that additional mechanisms besides direct BACE-1 inhibition also contribute to their A β lowering properties. We assessed the impact of flavonoids on NF κ B activation since we have previously identified that compounds inhibiting NF κ B activation can prevent the β -cleavage of APP and A β production [18]. We observed a strong correlation between the NF κ B inhibition potency of the flavonoids and their ability to inhibit A β production suggesting that the preponderant mechanism responsible for the A β lowering properties of the flavonoids is related to their effect on NF κ B activation. Interestingly, many flavonoids are known to inhibit tyrosine kinases and hence may regulate NF κ B activity via their inhibitory action on multiple tyrosine kinases upstream of NF κ B. NF κ B has been shown previously to regulate the production of A β by regulating the β -cleavage of APP [18]. Others studies have shown that NF κ B inhibition can directly regulate BACE-1 expression level [14, 30-32]. We effectively observed that apigenin, luteolin and quercetin inhibit BACE-1 transcription in human neuronal SHSY cells.

Conclusion:

Altogether our data suggest a multimodal mechanism of action of flavonoids towards A β production as the direct inhibition of BACE-1 activity by flavonoids appears to only marginally account for their A β lowering properties. We suggest that the A β lowering properties of flavonoids are mainly mediated via their effect on NF κ B signaling which in turn affects the regulation of BACE-1 expression.

Acknowledgements:

We would like to thank Diane and Robert Roskamp for their generosity in helping to make this work possible. We thank Dr. Michael Wolfe (Harvard Medical School, Boston, Massachusetts, USA) for providing the 7W CHO APP overexpressing cells.

Authors' contributions:

DP conceived of the study, developed the methodology for studying APP processing and NF κ B activity, performed BACE-1 activity measurements, the statistical analyzes and drafted the manuscript. VM performed the molecular docking simulations. GA realized the RTQ-PCR experiments. DB and NP performed the western-blot and ELISAs. CB contributed to the NF κ B luciferase activity measurements and helped in writing the manuscript. MM critically evaluated the manuscript. All authors read and approved the final manuscript.

Competing interest:

The authors declare that they have no competing interests.

References:

- [1] Selkoe DJ. *Behav Brain Res*. 2008 **192**:106 [PMID: 18359102]
- [2] Vassar R *et al. Science* 1999 **286**: 735 [PMID: 10531052]
- [3] Sinha S *et al. Nature* 1999 **402**: 537 [PMID: 10591214]
- [4] Yan R *et al. Nature* 1999 **402**: 533 [PMID: 10591213]
- [5] Cole SL & Vassar R. *J Biol Chem*. 2008 **283**: 29621 [PMID: 18650431]
- [6] Iwatsubo T. *Rinsho Shinkeigaku* 2004 **44**: 768 [PMID: 15651286]
- [7] Cai H *et al. Nat Neurosci*. 2001 **4**:233 [PMID: 11224536]
- [8] Dai Q *et al. Am J Med*. 2006 **119**: 751 [PMID: 16945610]
- [9] Commenges D *et al. Eur J Epidemiol*. 2000 **16**: 357 [PMID: 10959944]
- [10] Napryeyenko O *et al. Arzneimittelforschung* 2007 **57**: 4 [PMID: 17341003]
- [11] Mazza M *et al. Eur J Neurol*. 2006 **13**: 981 [PMID: 16930364]
- [12] Shimmyo Y *et al. Biochim Biophys Acta*. 2008 **1780**: 819 [PMID: 18295609]
- [13] Koo EH & Squazzo SL. *J Biol Chem*. 1994 **269**: 17386 [PMID: 8021238]
- [14] Paris D *et al. J Neuroinflammation*. 2010 **7**:17 [PMID: 20211007]

- [15] Ait-Ghezala G *et al.* *Brain Res Mol Brain Res.* 2005 **140**:73 [PMID: 16182406]
- [16] Garcia-Lafuente A *et al.* *Inflamm Res.* 2009 **58**: 537 [PMID: 19381780]
- [17] Ait-Ghezala G *et al.* *Eur J Neurosci.* 2007 **25**: 1685 [PMID: 17432959]
- [18] Paris D *et al.* *Neurosci Lett.* 2007 **415**: 11 [PMID: 17223266]
- [19] Bastianetto S *et al.* *Eur J Neurosci.* 2006 **23**: 55 [PMID: 16420415]
- [20] Iuvone T *et al.* *J Pharmacol Exp Ther.* 2006 **317**: 1143 [PMID: 16495207]
- [21] Kim JK *et al.* *Biosci Biotechnol Biochem.* 2010 **74**: 397 [PMID: 20139605]
- [22] Ono K *et al.* *J Neurochem.* 2003 **87**: 172 [PMID: 12969264]
- [23] Onozuka H *et al.* *J Pharmacol Exp Ther.* 2008 **326**: 739 [PMID: 18544674]
- [24] Rezai-Zadeh K *et al.* *J Cell Mol Med.* 2009 **13**: 574 [PMID: 18410522]
- [25] Choi YH *et al.* *Planta Med.* 2008 **74**: 1405 [PMID: 18666047]
- [26] Citron M. *J Neurosci Res.* 2002 **70**: 373 [PMID: 12391600]
- [27] Patel S *et al.* *J Mol Biol.* 2004 **343**: 407 [PMID: 15451669]
- [28] Moitessier N *et al.* *J Med Chem.* 2006 **49**: 5885 [PMID: 17004704]
- [29] Eriksson MA *et al.* *J Med Chem.* 1999 **42**: 868 [PMID: 10072684]
- [30] Wang S & Jia J. *Am J Med Genet B Neuropsychiatr Genet.* 2010 **153B**: 159 [PMID: 19441127]
- [31] Sambamurti K *et al.* *FASEB J.* 2004 **18**: 1034 [PMID: 15059975]
- [32] Buggia-Prevot V *et al.* *J Biol Chem.* 2008 **283**: 10037 [PMID: 18263584]

Edited by P Kanguane

Citation: Paris *et al.* *Bioinformation* 6(6): 229-236 (2011)
provided the original author and source are credited.

License statement: This is an open-access article, which permits unrestricted use, distribution, and reproduction in any medium, for non-commercial purposes,

Supplementary material:

Table 1: Summary of docking scores for different flavonoids using two different crystal structures for BACE-1 (2B8L and 2QMF). Compounds were ranked as a function of their IC50 for BACE-1 activity. A weak docking score was observed showing that flavonoids have a low affinity for the BACE-1 catalytic site.

Compound	pIC50	Glide XP Score (2B8L)	Glide XP Score (2QMF)	Ranking Based on pIC50	Ranking Based on 2B8L Docking	Ranking Based on 2QMF
Baicalein	-1	-5.34	-8.81	1	7	1
Kaemferol	-1.44	-8.42	-7.11	2	4	3
Quercetin	-1.48	-9.19	-8.04	3	2	2
Apigenin	-1.53	-9.01	-3.18	4	3	13
Luteolin	-1.75	-4.2	-5.37	5	10	6
Daizein	-1.91	-2.91	-4.66	6	12	9
Genistein	-1.94	-3.57	-5.23	7	11	8
PD98059	-2.24	-5.55	-3.79	8	6	10
β -naphthoflavone	-2.26	-4.77	-3.54	9	8	11
Taxifolin	-2.35	-10.49	-5.84	10	1	4
α -naphthoflavone	-2.83	-1.07	-3.44	11	13	12
Acacetin	-5.7	-4.27	-5.69	12	9	5
Aminogenistein	-5.7	-6.75	-5.36	13	5	7